

# THEORETICAL EVIDENCE IN SUPPORT OF THE EXTENDED ABELIAN RANK ONE STARK CONJECTURE FOR THE BASE FIELD $\mathbb{Q}$

DANIEL VALLIÈRES

ABSTRACT. The main goal of this paper is to study two infinite families of abelian extensions of  $\mathbb{Q}$ , one with Galois groups of the type  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z}$ , where  $m$  is an odd integer, and one with Galois groups of the type  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/\ell\mathbb{Z} \times \mathbb{Z}/\ell\mathbb{Z}$ , where  $\ell$  is an odd prime. We show partial results towards the extended abelian rank one Stark conjecture for the first family and we show that the extended abelian rank one Stark conjecture is true for the second one.

## CONTENTS

1. Introduction	1
Acknowledgement	2
1.1. Notation	2
1.2. Background	2
2. A generalization of the Brumer-Stark conjecture	3
2.1. Local version of the extension of the Brumer-Stark conjecture	5
2.2. Further remarks	8
2.3. Non semi-simplicity	9
3. Two infinite families of abelian extensions of $\mathbb{Q}$	10
3.1. The first family of abelian extensions of $\mathbb{Q}$ to be studied	10
The ramified prime $p$ .	11
The unramified prime $\ell$ .	11
3.2. The second family of abelian extensions of $\mathbb{Q}$ to be studied	12
The ramified primes $p_1$ and $p_2$ .	13
The unramified primes $p_3, \dots, p_{\ell+1}$ .	13
4. Conclusion	16
References	16

## 1. INTRODUCTION

The extended abelian rank one Stark conjecture was stated for the first time in [3]. Our approach is based on [15] and we refer to this latter paper for the statement of the conjecture (Conjecture 3.6). See also [2] where arbitrary orders of vanishing are treated. Here, we work exclusively with the  $S$ -version of the conjecture. For a given 1-cover  $S$  (see §1.2 below for the definitions of a 1-cover and  $S_{min}$ ), rather than studying the extended abelian rank one Stark conjecture, we study Question 4.2 of [15] (denoted by  $St(K/k, S, v)$  for a given 1-cover  $S$  and a place  $v \in S_{min}$ ). It was shown that an affirmative answer to  $St(K/k, S, v)$  for all  $v \in S_{min}$  implies the extended abelian rank one Stark conjecture (see Proposition 4.3 of [15]).

In this paper, we will be concerned almost exclusively with the case where  $v \in S_{min}$  is a finite prime. In this case, as explained in [15], it is possible to formulate an extension of the classical Brumer-Stark conjecture which, if true, implies that  $St(K/k, S, v)$  has an affirmative answer. This extension of the

---

*Date:* July 22, 2012.

This research was possible thanks to a research grant from the DAAD.

Brumer-Stark conjecture is recalled in §2 (Question 2.1). Question 2.1 will be the main object of study of this paper. In §1.1, we fix some of our basic notation and in §1.2, we recall some needed background information regarding the extended abelian rank one Stark conjecture. In §2.1, we formulate local versions of Question 2.1 and we prove some reduction statements similar as the ones contained in [6] for the classical Brumer-Stark conjecture. In §3, we use the previous results of §2.1 and some previous work of Greither and Kučera [4] and of Smith [12] in order to study two infinite families of abelian extensions of  $\mathbb{Q}$ .

**Acknowledgement.** The author would like to thank Cornelius Greither for suggesting to look at the example presented in §3.2. He would also like to thank Cornelius Greither, Radan Kučera and Barry Smith for explaining to him some of their previous results.

**1.1. Notation.** Given a finite abelian extension  $K/k$  of number fields with Galois group  $G$ , the symbol  $S(K/k)$  denotes the set of primes in  $k$  which are either archimedean or ramified in  $K/k$ . If  $S$  is any finite set of places of  $k$  containing  $S(K/k)$ , we denote the  $S$ -equivariant  $L$ -function by  $\theta_{K/k,S}(s)$ . When necessary, we shall write

$$\theta_{K/k,S}(0) = \theta_{K/k,S(K/k)}(0) \cdot \omega_{K/k,S},$$

where

$$\omega_{K/k,S} = \prod_{\mathfrak{p} \in S \setminus S(K/k)} (1 - \sigma_{\mathfrak{p}}^{-1}),$$

and  $\sigma_{\mathfrak{p}}$  denotes the Frobenius automorphism associated to the unramified prime  $\mathfrak{p}$  in  $K/k$ . In order to simplify the notation, we write  $\theta_{K/k}(0)$  instead of  $\theta_{K/k,S(K/k)}(0)$  (i.e. we drop the subscript  $S$  if  $S$  is minimal).

If the top field  $K$  is a  $CM$ -field, we denote the complex conjugation in  $G$  by  $j$ .

The subgroup of  $K^\times$  consisting of anti-units will be denoted by  $K^0$ . We remind the reader that  $\alpha \in K^\times$  is an anti-unit if  $|\alpha|_w = 1$  for all archimedean places  $w$  of  $K$ . We use the symbol  $w_K$  for the number of roots of unity in  $K^\times$ . If  $n | w_K$ , the subgroup of  $K^\times$  consisting of elements  $\lambda \in K^\times$  such that  $K(\lambda^{1/n})/k$  is an abelian extension of number fields will be denoted by  $\mathcal{A}(K/k, n)$ . This last property does not depend on the choice of  $\lambda^{1/n}$ . Both  $K^0$  and  $\mathcal{A}(K/k, n)$  are  $G$ -modules.

If  $A$  is a subgroup of  $I_K$ , the group of fractional ideals of  $K$ , and  $n | w_K$ , we define  $\text{BrSt}_{\mathbb{Z}[G]}^{(n)}(A)$  to be the set of elements  $\alpha \in \mathbb{Z}[G]$  for which, given any  $\mathfrak{a} \in A$ , there exists  $\eta(\mathfrak{a}) \in K^0 \cap \mathcal{A}(K/k, n)$  such that

$$\mathfrak{a}^\alpha = (\eta(\mathfrak{a})).$$

The set  $\text{BrSt}_{\mathbb{Z}[G]}^{(n)}(A)$  is an ideal of  $\mathbb{Z}[G]$ . The classical Brumer-Stark conjecture says

$$w_K \theta_{K/k,S}(0) \in \text{BrSt}_{\mathbb{Z}[G]}^{(w_K)}(I_K),$$

for any finite set  $S$  of places of  $k$  containing  $S(K/k)$ . It is known to be true for every finite abelian extension of  $\mathbb{Q}$ . See for instance [9].

If  $A$  is a finite abelian group, its  $p$ -Sylow subgroup will be denoted by  $A\{p\}$ .

**1.2. Background.** Let  $K/k$  be a finite abelian extension of number fields and  $S$  a finite set of places of  $k$  containing  $S(K/k)$ . One of the hypotheses of the classical abelian rank one Stark conjecture, as stated in [13], requires that there is a split prime in  $S$ . This condition implies that the order of vanishing of the  $S$ -imprimitive  $L$ -function  $L_{K/k,S}(s, \chi)$  is at least one for all non-trivial characters  $\chi$  of  $\text{Gal}(K/k)$ . This follows from the following theorem.

**Theorem 1.1.** *Let  $K/k$  be an abelian extension of number fields and  $S$  a finite set of primes of  $k$  containing the infinite ones. We have*

$$\text{ord}_{s=0} L_{K/k,S}(s, \chi) = \begin{cases} |S| - 1, & \text{if } \chi = \chi_1, \\ |\{v \in S \mid G_v \subseteq \text{Ker}(\chi)\}|, & \text{if } \chi \neq \chi_1, \end{cases}$$

where  $G_v$  denotes the decomposition group associated to the place  $v$  in  $G$ .

**Proof:**

See [14], page 24, Proposition 3.4.

Q.E.D.

On the other hand, it is not difficult to give examples of abelian extensions  $K/k$  and finite sets of places  $S$ , not containing any split prime, for which all the  $S$ -imprimitive  $L$ -functions vanish with order of vanishing at least one as well. Such a set  $S$  is called a 1-cover:

**Definition 1.2.** Let  $K/k$  be an abelian extension of number fields with Galois group  $G$  and let  $\Lambda$  be any subset of  $\widehat{G} = \text{Hom}_{\mathbb{Z}}(G, \mathbb{C}^{\times})$ . Let  $S$  be any finite set of primes of  $k$  (perhaps not containing the ramified nor the archimedean primes). The set  $S$  is said to be a 1-cover for  $\Lambda$  if the following two conditions hold:

- (1) For all non-trivial  $\chi \in \Lambda$ , there exists at least one prime  $v \in S$  such that  $G_v \subseteq \text{Ker}(\chi)$ ,
- (2) If the trivial character is in  $\Lambda$ , then  $|S| \geq 2$ .

In the case where  $\Lambda = \widehat{G}$ , we also say that  $S$  is a 1-cover for  $G$  (or for  $K/k$ ) rather than for  $\widehat{G}$ .

If  $S$  is a 1-cover for  $K/k$  containing  $S(K/k)$ , then  $\text{ord}_{s=0} L_{K/k,S}(s, \chi) \geq 1$  for all  $\chi \in \widehat{G}$  by Theorem 1.1. *Unless otherwise stated, we always suppose that a 1-cover  $S$  for  $K/k$  contains  $S(K/k)$ .* We need one more definition.

**Definition 1.3.** Let  $K/k$  be an abelian extension of number fields with Galois group  $G$  and  $S$  a 1-cover for  $K/k$ . The set of characters  $\chi \in \widehat{G}$  whose  $S$ -imprimitive  $L$ -functions have order of vanishing precisely one will be denoted by  $\widehat{G}_{1,S}$ . Moreover, the set  $S_{min}$  is defined as follows: It consists of all primes  $v \in S$  for which there exists  $\chi \in \widehat{G}_{1,S}$ ,  $\chi \neq \chi_1$ , such that  $G_v \subseteq \text{Ker}(\chi)$ .

In other words,  $S_{min}$  precisely consists of the places in  $S$  which are responsible for the vanishing of the  $S$ -imprimitive  $L$ -functions associated to non-trivial characters having order of vanishing exactly one.

It is natural to try to generalize the abelian rank one Stark conjecture to handle these more general sets  $S$  consisting of 1-covers. This is the purpose of the extended abelian rank one Stark conjecture of Erickson-Stark contained in [3].

In [15], the present author formulated and studied a stronger Question involving one  $v \in S_{min}$  at the time (this Question is denoted by  $St(K/k, S, v)$  where  $S$  is a 1-cover and  $v \in S_{min}$ ). It was shown that if  $St(K/k, S, v)$  has an affirmative answer for all  $v \in S_{min}$ , then the extended abelian rank one Stark conjecture is true. This Question seems to be easier to study both theoretically and numerically.

If  $v \in S_{min}$  is a *finite* prime, the present author also formulated in [15] a generalization of the Brumer-Stark conjecture which in the case of a positive answer implies that  $St(K/k, S, v)$  also has an affirmative answer. This generalization of the Brumer-Stark conjecture is the main object of study of this paper. We recall its precise statement in the next section.

## 2. A GENERALIZATION OF THE BRUMER-STARK CONJECTURE

The setting is as follows: Let  $K/k$  be a finite abelian extension of number fields and  $S$  a 1-cover for  $K/k$ . Suppose that  $|S| \geq 3$  and  $S \neq S_{min}$ . Let  $\mathfrak{p} \in S_{min}$  be a finite prime. As in [15], let  $L^{(\mathfrak{p})} = K^{G_{\mathfrak{p}}}$ ,  $\Gamma_{\mathfrak{p}} = \text{Gal}(L^{(\mathfrak{p})}/k)$ ,  $n_{\mathfrak{p}} = |G_{\mathfrak{p}}|$  and  $R_{\mathfrak{p}} = S \setminus \{\mathfrak{p}\}$ . If  $\mathfrak{p} \in S_{min}$  is fixed, then we write  $L, \Gamma, R, n$  instead of  $L^{(\mathfrak{p})}, \Gamma_{\mathfrak{p}}, R_{\mathfrak{p}}, n_{\mathfrak{p}}$ , and similarly for other notations depending on  $\mathfrak{p} \in S_{min}$ .

- (1) If  $\mathfrak{p} \in S_{min}$  is ramified in  $K/k$  define the subgroup  $A = A_{\mathfrak{p}}$  of  $I_L$  to be the subgroup generated by the primes  $\mathfrak{P}$  of  $L$  lying above  $\mathfrak{p}$ .
- (2) If  $\mathfrak{p} \in S_{min}$  is unramified in  $K/k$ , let  $A = A_{\mathfrak{p}}$  be the subgroup of  $I_L$  generated by the primes  $\mathfrak{Q}$  of  $L$  lying above primes  $\mathfrak{q}$  of  $k$  satisfying

$$\left( \frac{K/k}{\mathfrak{q}} \right) = \left( \frac{K/k}{\mathfrak{p}} \right).$$

The extension of the Brumer-Stark conjecture alluded to above is the following (Question 4.7 of [15]).

**Question 2.1** (Extension of the Brumer-Stark conjecture). *Let  $\mathfrak{p} \in S_{min}$  be a finite prime. In the setting as above, do we have*

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \mathbb{Z}[\Gamma]?$$

Moreover, do we have

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L)}(A),$$

where  $A$  is the subgroup of  $I_L$  defined above?

In all theoretical and numerical examples that have been investigated so far, one can in fact take  $I_L$  instead of  $A$  in the previous Question. Numerical examples have been computed only in the case where the base field is  $\mathbb{Q}$ , see [15] for some examples and [16] for more. It would be interesting to do computations with other base fields than  $\mathbb{Q}$ .

We remark that since we assume  $\mathfrak{p} \in S_{min}$  is a finite prime, the field  $k$  is necessarily totally real and  $K$  totally complex. Now, since  $K/k$  is Galois and  $k$  is totally real, every complex embedding of  $K$  induces a complex conjugation. If  $w$  is a (necessarily complex) place of  $K$ , we denote the corresponding complex conjugation by  $\sigma_w$ . The fact that  $K/k$  is abelian implies that  $\sigma_w$  depends only on the place  $v$  of  $k$  lying below  $w$ . Thus, it will be denoted by  $\sigma_v$  rather than  $\sigma_w$ . By the decomposition group associated to the infinite place  $v$  of  $k$ , we mean, as usual,  $G_v := \langle \sigma_v \rangle$ . The field  $L$  also has to be totally complex, otherwise  $G_v \subseteq G_{\mathfrak{p}}$  for some real place  $v$  of  $k$ , and this is impossible, since  $\mathfrak{p}$  is assumed to be in  $S_{min}$ . If we are in the situation where  $K$  is a  $CM$ -field, then  $L$  is also a  $CM$ -field as subfields of  $CM$ -fields are either totally real or  $CM$ -fields.

As another remark, we note that if  $R$  is any finite set of primes containing  $S(K/k)$  and  $\mathfrak{p}$  is a prime of  $k$  which splits completely in  $K/k$ , then  $S = R \cup \{\mathfrak{p}\}$  is automatically a 1-cover. There are only two possibilities: Either  $S_{min}$  is empty or  $S_{min} = \{\mathfrak{p}\}$ . In the former case,  $\theta_{K/k,R}(0) = 0$  and there is nothing to show, whereas in the latter case  $K = L$ ,  $n = 1$  and  $A = I_K^{spl}$  is the group of fractional ideals generated by the primes  $\mathfrak{P}$  of  $K$  lying above primes  $\mathfrak{p}$  of  $k$  splitting completely in  $K/k$ . Since it is known (see the classical book on the subject [14] or Lemma 2.3 below) that

$$w_K \theta_{K/k,R}(0) \in \text{BrSt}_{\mathbb{Z}[G]}^{(w_K)}(P_K),$$

where  $P_K$  is the group of principal ideals of  $K$  and  $I_K$  is generated by  $P_K$  and  $I_K^{spl}$ , we get back the original Brumer-Stark conjecture.

We will refer to the following hypothesis as the Integrality Property.

**Hypothesis 2.2** (Integrality Property). *One has*

$$\text{Ann}_{\mathbb{Z}[\Gamma]}(\mu_L) \cdot \frac{\theta_{L/k,R}(0)}{n} \subseteq \mathbb{Z}[\Gamma].$$

If the Integrality Property is satisfied, then the first part of Question 2.1 holds true, since  $w_L \in \text{Ann}_{\mathbb{Z}[\Gamma]}(\mu_L)$ . It was shown in [15] (Theorem 4.30) that if  $\mathfrak{p} \in S_{min}$  is unramified in  $K/k$ , then the Integrality Property follows from a conjecture of Gross (Conjecture 7.6 of [7]). This case of the Gross conjecture is known to be true, see for instance Lemma 2.3 in [8]. Therefore, the Integrality Property is true if the finite prime  $\mathfrak{p} \in S_{min}$  is unramified in  $K/k$ . If the finite prime  $\mathfrak{p} \in S_{min}$  is ramified in  $K/k$ , then we cannot say anything in theory, but the numerical computations agree with it so far.

**Lemma 2.3.** *Suppose that the Integrality Property holds true. Then*

$$(L^\times)^{\frac{w_L \cdot \theta_{L/k,R}(0)}{n}} \subseteq L^0 \cap \mathcal{A}(L/k, w_L).$$

**Proof:**

This follows from Lemmas 4.10 and 4.12 of [15].

Q.E.D.

**2.1. Local version of the extension of the Brumer-Stark conjecture.** This section (§2.1) is based on [6] where the corresponding results are proved for the classical Brumer-Stark conjecture. The proofs are similar and we include them here only for the sake of completeness relative to the setting of Question 2.1.

Suppose that  $m \mid w_L$ . If the Integrality Property holds true then it makes sense to say

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(m)}(A),$$

where  $A$  is now a subgroup of  $Cl_L$  rather than  $I_L$ . Indeed, if  $\mathfrak{a}$  is a fractional ideal such that there exists  $\eta(\mathfrak{a}) \in L^\times$  satisfying

- (1)  $\mathfrak{a}^{\frac{w_L \cdot \theta_{L/k,R}(0)}{n}} = (\eta(\mathfrak{a}))$ ,
- (2)  $|\eta(\mathfrak{a})|_w = 1$  for all infinite places  $w$  of  $L$ ,
- (3) The extension  $L(\eta(\mathfrak{a})^{1/m})/k$  is abelian,

and  $\mathfrak{b} = \lambda \cdot \mathfrak{a}$  for some  $\lambda \in L^\times$ , then  $\mathfrak{b}$  satisfies the same three properties with  $\eta(\mathfrak{b}) = \lambda^{\frac{w_L \theta_{L/k,R}(0)}{n}} \cdot \eta(\mathfrak{a})$  by Lemma 2.3.

**Proposition 2.4.** *Suppose the Integrality Property holds true and let  $A$  be any subgroup of  $Cl_L$ . Let  $p$  be any prime number and set  $w_{L,p} = |w_L|_p^{-1}$ , i.e.  $w_{L,p}$  is the exact power of  $p$  dividing  $w_L$ . One has*

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L)}(A\{p\})$$

if and only if

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_{L,p})}(A\{p\}).$$

**Proof:**

One direction is clear. So suppose that

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_{L,p})}(A\{p\}),$$

and write  $w_L = m \cdot w_{L,p}$  for some integer  $m$  relatively prime to  $p$ . Since  $(m, p) = 1$ , the group morphism

$$A\{p\} \longrightarrow A\{p\},$$

defined by  $[\mathfrak{a}] \mapsto [\mathfrak{a}]^m$  is an isomorphism. Let  $[\mathfrak{a}] \in A\{p\}$  and let  $[\mathfrak{b}] \in A\{p\}$  be such that  $[\mathfrak{b}]^m = [\mathfrak{a}]$ . By hypothesis, there exists  $\eta(\mathfrak{b}) \in L^0 \cap \mathcal{A}(L/k, w_{L,p})$  such that

$$\mathfrak{b}^{\frac{w_L \theta_{L/k,R}(0)}{n}} = (\eta(\mathfrak{b})).$$

Now, there exists  $\lambda \in L^\times$  such that  $\mathfrak{a} = \lambda \mathfrak{b}^m$ . We then have

$$\mathfrak{a}^{\frac{w_L \theta_{L/k,R}(0)}{n}} = \left( \lambda^{\frac{w_L \theta_{L/k,R}(0)}{n}} \eta(\mathfrak{b})^m \right).$$

By Lemma 2.3,

$$\lambda^{\frac{w_L \theta_{L/k,R}(0)}{n}} \in L^0 \cap \mathcal{A}(L/k, w_L),$$

and since  $\eta(\mathfrak{b}) \in L^0 \cap \mathcal{A}(L/k, w_{L,p})$ , we conclude that  $\eta(\mathfrak{b})^m \in L^0 \cap \mathcal{A}(L/k, w_L)$ . This is what we wanted to show.

Q.E.D.

This last proposition allows one to study Question 2.1 one prime at a time:

**Corollary 2.5.** *If  $p$  is a prime number, let  $n_p = |n|_p^{-1}$ . The following statements are equivalent:*

- (1)  $\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L)}(A)$ ,
- (2)  $\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_{L,p})}(A\{p\})$ , for all prime numbers  $p$ .

$$(3) \quad \frac{w_L \cdot \theta_{L/k,R}(0)}{n_p} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L,p)}(A\{p\}), \text{ for all prime numbers } p.$$

**Proof:**

The equivalence between (1) and (2) follows directly from Proposition 2.4. The proof of the equivalence between (2) and (3) is similar and left to the reader.

Q.E.D.

The next proposition shows that assuming the usual Brumer-Stark conjecture, Question 2.1 does not say anything new at primes  $p$  which are relatively prime to  $n$ .

**Proposition 2.6.** *Suppose that the Integrality Property holds true and also that the usual Brumer-Stark conjecture is true for  $L/k$ . If  $p$  is a prime satisfying  $(p, n) = 1$ , then*

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L,p)}(Cl_L\{p\}).$$

**Proof:**

Let  $[\mathfrak{a}] \in Cl_L\{p\}$  and  $p^e$  be the exponent of  $Cl_L\{p\}$ . Let  $\lambda_1 \in K^\times$  be such that  $\mathfrak{a}^{p^e} = (\lambda_1)$ . Since  $(p, n) = 1$ , there exist  $s, t \in \mathbb{Z}$  such that  $sp^e + tn = 1$ . Therefore,

$$\begin{aligned} \mathfrak{a}^{\frac{w_L \theta_{L/k,R}(0)}{n}} &= \mathfrak{a}^{s \cdot p^e \cdot \frac{w_L \theta_{L/k,R}(0)}{n}} \cdot \mathfrak{a}^{t \cdot w_L \theta_{L/k,R}(0)} \\ &= \left( \lambda_1^{s \cdot \frac{w_L \theta_{L/k,R}(0)}{n}} \right) \cdot \mathfrak{a}^{t \cdot w_L \theta_{L/k,R}(0)}. \end{aligned}$$

By Lemma 2.3, we have  $\lambda_1^{\frac{w_L \theta_{L/k,R}(0)}{n}} \in L^0 \cap \mathcal{A}(L/k, w_L)$ . Moreover, the usual Brumer-Stark conjecture for the special value  $w_L \theta_{L/k,R}(0)$  shows that the ideal  $\mathfrak{a}^{w_L \theta_{L/k,R}(0)}$  is principal generated by an element in  $L^0 \cap \mathcal{A}(L/k, w_L)$ . This is what we wanted to show.

Q.E.D.

Hence, we just have to focus on the primes  $p$  dividing  $n$ .

**Proposition 2.7.** *Let  $A$  be any subgroup of  $Cl_L$  and suppose that the Integrality Property holds true. Let  $p$  be a prime number dividing  $n$ . Suppose moreover that  $(p, w_L) = 1$ . Then*

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L,p)}(A\{p\})$$

if and only if

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{Ann}_{\mathbb{Z}[\Gamma]}(A\{p\}).$$

**Proof:**

Again, one direction is clear, so suppose that

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{Ann}_{\mathbb{Z}[\Gamma]}(A\{p\}),$$

and let  $[\mathfrak{a}] \in A\{p\}$ . Let  $p^e$  be the exponent of  $A\{p\}$  and let  $\lambda \in L^\times$  be such that  $\mathfrak{a}^{p^e} = (\lambda)$ . Since  $(p, w_L) = 1$ , there exist  $s, t \in \mathbb{Z}$  such that  $sp^e + tw_L = 1$ . By hypothesis, there exists  $\eta(\mathfrak{a}) \in L^\times$  such that

$$\mathfrak{a}^{\frac{w_L \cdot \theta_{L/k,R}(0)}{n}} = (\eta(\mathfrak{a})),$$

hence we have

$$\mathfrak{a}^{\frac{w_L \cdot \theta_{L/k,R}(0)}{n}} = \left( \lambda^{s \cdot \frac{w_L \cdot \theta_{L/k,R}(0)}{n}} \eta(\mathfrak{a})^{tw_L} \right).$$

Setting

$$\eta'(\mathfrak{a}) = \lambda^{s \cdot \frac{w_L \cdot \theta_{L/k,R}(0)}{n}} \eta(\mathfrak{a})^{tw_L},$$

we see that  $\eta'(\mathfrak{a}) \in \mathcal{A}(L/k, w_L)$  by Lemma 2.3.

As explained after Question 2.1, the field  $L$  is necessarily totally complex. For every (necessarily complex) place  $w$  of  $L$ , let  $\sigma_w$  be the associated complex conjugation. From Lemma 4.10 in [15], we have

$$(1 + \sigma_w) \frac{w_L \theta_{L/k, R}(0)}{n} = 0,$$

for all infinite places  $w$  of  $L$ . Since  $1 - \sigma_w = 2 - (1 + \sigma_w)$ , we get

$$(1 - \sigma_w) \frac{w_L \theta_{L/k, R}(0)}{n} = 2 \cdot \frac{w_L \theta_{L/k, R}(0)}{n}.$$

Hence, letting  $d = [L : \mathbb{Q}]/2$ , we have

$$\prod_w (1 - \sigma_w) \cdot \frac{w_L \theta_{L/k, R}(0)}{n} = 2^d \frac{w_L \theta_{L/k, R}(0)}{n},$$

where the product is over all (necessarily complex) infinite places of  $L$ . Since  $(p, w_L) = 1$  and  $w_L$  is even, we see that there exists  $s, t \in \mathbb{Z}$  such that  $s2^d + tp^e = 1$ . Then

$$\begin{aligned} \mathfrak{a}^{\frac{w_L \theta_{L/k, R}(0)}{n}} &= \left( \lambda^{\frac{tw_L \theta_{L/k, R}(0)}{n}} \right) \mathfrak{a}^{\frac{s2^d w_L \theta_{L/k, R}(0)}{n}} \\ &= \left( \lambda^{\frac{tw_L \theta_{L/k, R}(0)}{n}} \eta'(\mathfrak{a})^s \Pi_w(1 - \sigma_w) \right). \end{aligned}$$

Letting

$$\eta''(\mathfrak{a}) = \lambda^{\frac{tw_L \theta_{L/k, R}(0)}{n}} \eta'(\mathfrak{a})^s \Pi_w(1 - \sigma_w),$$

we see that  $\eta''(\mathfrak{a}) \in L^0 \cap \mathcal{A}(L/k, w_L)$ .

Q.E.D.

If  $A$  is a subgroup of the group of fractional ideals  $I_L$  and  $m \mid w_L$ , we define  $\text{Ab}_{\mathbb{Z}[\Gamma]}^{(m)}(A)$  to be the set of elements  $\alpha \in \mathbb{Z}[\Gamma]$  such that there exists  $\eta(\mathfrak{a}) \in \mathcal{A}(L/k, m)$  satisfying

$$\mathfrak{a}^\alpha = (\eta(\mathfrak{a})).$$

In other words, we are just dropping the anti-unit condition. The set  $\text{Ab}_{\mathbb{Z}[\Gamma]}^{(m)}(A)$  is an ideal of  $\mathbb{Z}[\Gamma]$  and we clearly have the inclusion

$$\text{BrSt}_{\mathbb{Z}[\Gamma]}^{(m)}(A) \subseteq \text{Ab}_{\mathbb{Z}[\Gamma]}^{(m)}(A).$$

Again, if the Integrality Property holds true, Lemma 2.3 implies that it makes sense to say

$$\frac{w_L \cdot \theta_{L/k, R}(0)}{n} \in \text{Ab}_{\mathbb{Z}[\Gamma]}^{(m)}(A)$$

if now  $A$  is a subgroup of  $Cl_L$  rather than  $I_L$ .

**Proposition 2.8.** *Let  $A$  be any subgroup of  $Cl_L$  and suppose that the Integrality Property holds true. Let  $p$  be an odd prime number dividing  $n$ . Suppose moreover that  $p \mid w_L$ . Then*

$$\frac{w_L \cdot \theta_{L/k, R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L, p)}(A\{p\})$$

if and only if

$$\frac{w_L \cdot \theta_{L/k, R}(0)}{n} \in \text{Ab}_{\mathbb{Z}[\Gamma]}^{(w_L, p)}(A\{p\}).$$

**Proof:**

The proof is similar to the second part of the proof of Proposition 2.7 and is left to the reader.

Q.E.D.

We can summarize the previous discussion as follows always assuming the Integrality Property. If  $p$  is a prime number and  $(p, n) = 1$ , we have nothing to do assuming the usual Brumer-Stark conjecture. Suppose now  $p \mid n$ . In order to give an affirmative answer to Question 2.1, then

- (1) If  $(p, w_L) = 1$ , we just have to show an annihilation statement.

- (2) If  $p \mid w_L$  and  $p$  is odd, we have to show an annihilation statement and the abelian condition.
- (3) If  $p = 2$ , then we have to show an annihilation statement, the abelian condition and the anti-unit part.

Suppose that  $L$  is a  $CM$ -field and  $p$  is an odd prime. Assuming the usual Brumer-Stark conjecture, we just have to deal with the minus part of the class group as the following proposition shows.

**Proposition 2.9.** *Suppose that the Integrality Property holds true and that the usual Brumer-Stark conjecture is true for  $L/k$ . Suppose also that  $L$  is a  $CM$ -field and  $p$  is an odd prime dividing  $n$ . Letting  $n_p = |n|_p^{-1}$ , we have*

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n_p} \in \text{BrSt}_{\mathbb{Z}[G]}^{(w_L)}(Cl_L\{p\}^+).$$

**Proof:**

If  $[\mathfrak{a}] \in Cl_L\{p\}^+$ , then  $\mathfrak{a}^{1-j} = (\lambda_1)$  for some  $\lambda_1 \in L^\times$ . Moreover, we have the following identity

$$(1-j) \frac{w_L \cdot \theta_{L/k,R}(0)}{n_p} = 2 \cdot \frac{w_L \cdot \theta_{L/k,R}(0)}{n_p},$$

since  $\theta_{L/k,R}(0) \in \mathbb{Q}[G]^-$ . Since  $(2, n_p) = 1$ , there exist  $s, t \in \mathbb{Z}$  such that  $s2 + tn_p = 1$ . Thus,

$$\mathfrak{a}^{\frac{w_L \theta_{L/k,R}(0)}{n_p}} = \mathfrak{a}^{s \cdot 2 \cdot \frac{w_L \theta_{L/k,R}(0)}{n_p}} \cdot \mathfrak{a}^{t \cdot w_L \theta_{L/k,R}(0)}.$$

Now, we have

$$\mathfrak{a}^{2 \cdot \frac{w_L \theta_{L/k,R}(0)}{n_p}} = \mathfrak{a}^{(1-j) \cdot \frac{w_L \theta_{L/k,R}(0)}{n_p}} = \left( \lambda_1^{\frac{w_L \theta_{L/k,R}(0)}{n_p}} \right),$$

and  $\lambda_1^{\frac{w_L \theta_{L/k,R}(0)}{n_p}} \in L^0 \cap \mathcal{A}(L/k, w_L)$  by Lemma 2.3.

Moreover, the usual Brumer-Stark conjecture for the special value  $w_L \theta_{L/k,R}(0)$  shows that the ideal  $\mathfrak{a}^{w_L \theta_{L/k,R}(0)}$  is principal generated by an element in  $L^0 \cap \mathcal{A}(L/k, w_L)$ . This is what we wanted to prove.

Q.E.D.

**2.2. Further remarks.** It seems clear now that a stronger Brumer-Stark type statement for a certain subgroup  $A$  of  $I_L$  would imply a big part of the extended abelian rank one Stark conjecture. As explained at the beginning of §2, depending on whether  $\mathfrak{p}$  is ramified or not in  $K/k$ , we proposed two such subgroups which would be enough to have consequences for the extended abelian rank one Stark conjecture.

On the other hand, surprisingly perhaps, in all theoretical and numerical results so far, Question 2.1 holds true with  $A = I_L$ . The simplest possible type of fields for which Question 2.1 does not reduce to the classical Brumer-Stark conjecture is when  $K/k$  is biquadratic.

**Theorem 2.10.** *Suppose that  $K/k$  is a biquadratic extension of number fields with  $K$  totally complex and  $k$  totally real. Suppose also that  $S$  is a 1-cover which do not contain a split prime,  $|S| \geq 3$  and  $S \neq S_{\min}$ . Let  $\mathfrak{p} \in S_{\min}$  be a finite prime,  $L = K^{G_{\mathfrak{p}}}$ ,  $\Gamma = \text{Gal}(L/k)$  and  $R = S \setminus \{\mathfrak{p}\}$ . Note that  $|G_{\mathfrak{p}}| = 2$ . Then*

$$\frac{w_L \theta_{L/k,R}(0)}{2} \in \mathbb{Z}[\Gamma],$$

and moreover

$$\frac{w_L \theta_{L/k,R}(0)}{2} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L)}(I_L).$$

**Proof:**

Proposition 5.2 of [10] implies the following result for quadratic extensions. If  $M/k$  is a quadratic extension with Galois group  $\Gamma$  such that  $|S(M/k)| \geq 3$ , then

$$\frac{w_M \theta_{M/k}(0)}{2} \in \mathbb{Z}[\Gamma],$$

and

$$\frac{w_M \theta_{M/k}(0)}{2} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_M)}(I_M).$$

If we come back to our biquadratic extension  $K/k$  and subfield  $L$  (which is quadratic over  $k$  and also a  $CM$ -field), there are only two cases not covered by this result:

- (1) The base field  $k$  is  $\mathbb{Q}$  and only one finite prime ramifies in  $L/\mathbb{Q}$ .
- (2) The base field  $k$  is a real quadratic number field and the extension  $L/k$  is unramified.

As in the proof of Theorem 4.26 of [15], since  $S$  is a 1-cover and  $S \neq S_{min}$ , we know that  $|S| \geq 4$ . So in both cases, there is a prime  $\mathfrak{q} \in R$  which is unramified in  $L/k$ . Since  $\mathfrak{p} \in S_{min}$ , the Frobenius automorphism  $\sigma_{\mathfrak{q}}$  has to be the complex conjugation  $j \in \Gamma$ . Thus,

$$\begin{aligned} w_L \theta_{L/k,R}(0) &= w_L \cdot \theta_{L/k}(0) \cdot (1 - j) \cdot \beta \text{ for some } \beta \in \mathbb{Z}[\Gamma] \\ &= 2 \cdot w_L \cdot \theta_{L/k}(0) \cdot \beta, \end{aligned}$$

and hence

$$\frac{w_L \theta_{L/k,R}(0)}{2} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L)}(I_L).$$

This ends the proof, since the usual Brumer-Stark conjecture is known to be true for quadratic extensions.

Q.E.D.

It might be worthwhile to reinterpret a theorem of Erickson in terms of Question 2.1.

**Theorem 2.11** (Erickson). *Let  $S$  be a 1-cover for  $K/k$ . Moreover, suppose there exists a subset  $S' \subseteq S$  which consists only of unramified finite primes and such that  $S'$  is a 1-cover for  $K/k$ . Then, for every  $\mathfrak{p} \in S_{min}$ , one has*

$$\omega_{L/k,R} \in n \cdot \mathbb{Z}[\Gamma].$$

Therefore, there exists  $\alpha \in \mathbb{Z}[\Gamma]$  (which depends on  $\mathfrak{p}$ ) such that

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} = w_L \cdot \theta_{L/k}(0) \cdot \alpha,$$

and if the usual Brumer-Stark conjecture is true for  $L/k$ , then

$$\frac{w_L \cdot \theta_{L/k,R}(0)}{n} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L)}(I_L).$$

**Proof:**

See the proof of Theorem 6.1 in [3].

Q.E.D.

Let us remark that if this theorem applies, then  $S_{min}$  consists only of finite unramified primes.

**2.3. Non semi-simplicity.** Question 2.1 involves a “non semi-simple” situation as we now explain. In [15], we made the following definition.

**Definition 2.12.** Let  $G$  be a finite abelian group. A subgroup  $H$  is called *cocyclic* if  $G/H$  is a cyclic group. Moreover, if  $H$  is cocyclic and if  $K \subsetneq H$  implies that  $G/K$  is not cyclic, then we say that  $H$  is a *minimal cocyclic subgroup*.

We refer to §3.3 of [15] for the relation between minimal cocyclic subgroups and 1-covers.

**Lemma 2.13.** *Let  $G$  be a finite abelian group and let  $H$  be a subgroup. Then  $H$  is a minimal cocyclic subgroup of  $G$  if and only if  $H\{p\}$  is a minimal cocyclic subgroup of  $G\{p\}$  for all prime  $p$ .*

**Proof:**

This is clear using the fact that

$$G/H \simeq \bigoplus_p G\{p\}/H\{p\},$$

where the direct sum is over all prime numbers  $p$ .

Q.E.D.

Let us now come back to the setting of Question 2.1 for some finite prime  $\mathfrak{p} \in S_{min}$ .

**Proposition 2.14.** *Let  $p$  be a prime number. If  $p \mid n$ , then  $p \mid [L : k]$ .*

**Proof:**

Since  $\mathfrak{p} \in S_{min}$ , it implies that  $G_{\mathfrak{p}}$  is contained in a minimal cocyclic subgroup  $H$  of  $G$  (Theorems 3.15 and 3.16 of [15]). If  $p \nmid [L : k]$ , then  $G_{\mathfrak{p}}\{p\} = G\{p\}$ . Thus

$$G\{p\} = G_{\mathfrak{p}}\{p\} \subseteq H\{p\} \subseteq G\{p\},$$

and we conclude that  $H\{p\} = G\{p\}$ . This is a contradiction with Lemma 2.13.

Q.E.D.

By Proposition 2.6, we know that if  $(p, n) = 1$ , Question 2.1 does not say anything new assuming the usual Brumer-Stark conjecture. Now, if  $p \mid n$ , the last proposition says that we are studying an annihilation statement

$$\frac{w_L \cdot \theta_{L/k, R}(0)}{n} \in \text{Ann}_{\mathbb{Z}[\Gamma]}(Cl_L\{p\}),$$

in the case where  $p \mid [L : k]$ . This is the so-called “non semi-simple” case.

**Remark.** *Suppose we are in the setting of the question  $St(K/k, S, v)$  of [15] where  $v \in S_{min}$  is a real infinite place of  $k$ . The same argument as above shows that we necessarily have  $2 \mid [L : k]$  where  $L = K^{G_v}$ .*

### 3. TWO INFINITE FAMILIES OF ABELIAN EXTENSIONS OF $\mathbb{Q}$

**3.1. The first family of abelian extensions of  $\mathbb{Q}$  to be studied.** The following example is taken from [2] and was studied numerically in some cases in [15] and [16]. Let  $p, q$  be two odd prime numbers satisfying

$$p \equiv 1 \pmod{4}, \quad q \equiv 3 \pmod{4}, \quad \text{and} \quad \left(\frac{p}{q}\right) = 1.$$

Let  $K' = \mathbb{Q}(\zeta_q)^{D_p}$ , where  $D_p$  is the decomposition group associated to  $p$  in  $\mathbb{Q}(\zeta_q)/\mathbb{Q}$ . We remark that  $\mathbb{Q}(\sqrt{-q}) \subseteq K'$ . Let  $K'^+$  be the maximal real subfield of  $K'$ . Let  $\ell$  be an odd prime number, different from  $p$  and  $q$  and satisfying

$$\left(\frac{K'/\mathbb{Q}}{\ell}\right) \neq 1, \quad \left(\frac{K'^+/\mathbb{Q}}{\ell}\right) = 1, \quad \text{and} \quad \left(\frac{p}{\ell}\right) = -1,$$

(i.e.  $\ell$  splits completely in  $K'^+/\mathbb{Q}$ , but does not split completely in  $K'/\mathbb{Q}$  nor in  $\mathbb{Q}(\sqrt{p})/\mathbb{Q}$ ). Let  $K = K'(\sqrt{p})$ , then the Galois group  $G = \text{Gal}(K/\mathbb{Q})$  is isomorphic to

$$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/m\mathbb{Z},$$

where  $m = (q-1)/2f_p$ , and  $f_p$  is the inertia index of  $p$  in  $\mathbb{Q}(\zeta_q)$ . Since  $q \equiv 3 \pmod{4}$  we have  $(m, 2) = 1$ . Note that  $|G_p| = |G_\ell| = 2$  and  $G_p, G_\infty, G_\ell$  are precisely the three subgroups of index 2 in  $G$ . By Theorem 3.15 and Theorem 3.16 of [15], we see that  $S = \{\infty, p, q, \ell\}$  is a 1-cover for  $K/\mathbb{Q}$  and  $S_{min} = \{\infty, p, \ell\}$ .

If  $m = 1$ , then we are in the setting of Theorem 2.10. Otherwise, the simplest thing that could happen is that  $m$  is an odd prime. Here is our main theorem for this family of abelian extensions of  $\mathbb{Q}$ .

**Theorem 3.1.** *We have:*

- (1) *For the ramified prime  $p \in S_{min}$ , one has*

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{2} \in \mathbb{Z}[\Gamma].$$

Moreover,

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{2} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L)}(I_L).$$

Hence, Question 2.1 is true for  $p \in S_{min}$ .

(2) For the unramified prime  $\ell \in S_{min}$ , one has

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{2} \in \mathbb{Z}[\Gamma].$$

Suppose moreover that  $m$  is an odd prime and that 2 is a primitive root modulo  $m$ . Then

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{2} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L)}(I_L).$$

**Proof:**

**The ramified prime  $p$ .** Recall that  $L = K^{G_p}$ ,  $\Gamma = \text{Gal}(L/\mathbb{Q})$ ,  $R = S \setminus \{p\}$ , and  $|G_p| = 2$ . Since

$$\omega_R = 1 - \sigma_\ell^{-1},$$

and  $\sigma_\ell = j$  in  $\Gamma$ , we have

$$\begin{aligned} w_L \cdot \theta_{L/\mathbb{Q}, R}(0) &= w_L \cdot \theta_{L/\mathbb{Q}}(0) \cdot (1 - j) \\ &= 2 \cdot w_L \cdot \theta_{L/\mathbb{Q}}(0). \end{aligned}$$

Therefore,

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{2} = w_L \cdot \theta_{L/\mathbb{Q}}(0) \in \mathbb{Z}[\Gamma],$$

and

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{2} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L)}(I_L),$$

because of the usual Brumer-Stark conjecture for the special value  $w_L \cdot \theta_{L/\mathbb{Q}}(0)$ . Hence, point (1) of Theorem 3.1 is now proved.

**The unramified prime  $\ell$ .** Recall here that  $L = K^{G_\ell}$ ,  $\Gamma = \text{Gal}(L/\mathbb{Q})$ ,  $R = S \setminus \{\ell\}$ , and  $|G_\ell| = 2$ . As noted after Hypothesis 2.2, we have

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{2} \in \mathbb{Z}[\Gamma],$$

since  $\ell$  is unramified in  $K/\mathbb{Q}$ . Because of Corollary 2.5, Proposition 2.6 and noting that  $w_{L,2} = w_L = 2$ , we just have to show that

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{2} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(2)}(Cl_L\{2\}).$$

If  $m$  is an odd prime and 2 is a primitive root modulo  $m$ , then this follows from Theorem 5.3 of [12]. (In his notation, we have  $K_1 = L$ ,  $K_0 = \mathbb{Q}(\sqrt{-pq})$ ,  $k_1 = (K^+)^+$ ,  $k_0 = \mathbb{Q}$ , his  $S$  is our  $R$ ,  $S_1 = \{\infty, p\}$ , and  $r = 1$ .)

Q.E.D.

**Remark.** In [12], it was shown that if a certain Equation (7) were true, then the conclusion of point (2) of Theorem 3.1 would hold true if  $m$  is any odd prime. In other words, we would not need the hypothesis that 2 is a primitive root modulo  $m$ . See Proposition 5.2 of [12].

As for the extended abelian rank one Stark conjecture for this example, here is what we can say. In the case where  $m$  is an odd prime and 2 is a primitive root modulo  $m$ , Theorem 3.1 implies that  $St(K/k, S, p)$  and  $St(K/k, S, \ell)$  have an affirmative answer (Propositions 4.4 and 4.9 of [15]). We only have to study what is happening at  $\infty \in S_{min}$ , i.e. to study  $St(K/k, S, \infty)$ . Let now  $L = K^+ = K^{G_\infty}$ ,  $\Gamma = \text{Gal}(L/\mathbb{Q})$  and fix a place  $w$  of  $L$  lying above  $\infty$ . Assuming the Gross conjecture (Conjecture 7.6 of [7]) in the case where the split prime is the infinite one  $\infty$ , Theorem 4.27 of [15] implies the existence of an  $\eta \in L^0$  such that

$$\theta'_{L/\mathbb{Q}, S}(0) = - \sum_{\gamma \in \Gamma} \log |\eta^\gamma|_w \cdot \gamma^{-1}.$$

This case of the Gross conjecture when the base field is  $\mathbb{Q}$  is known to be true because of classical results (see §1.1 of [1] combined with the usual functorial properties of the Gross conjecture as top

change and enlarging the set  $S$ ). Because of Proposition 3.8 of [15], the extended abelian rank one Stark conjecture for  $K/k$  and  $S$  is true if and only if  $\eta \in \mathcal{A}(L/\mathbb{Q}, 2)$ , i.e. if the abelian condition of  $St(K/k, S, \infty)$  holds true.

**3.2. The second family of abelian extensions of  $\mathbb{Q}$  to be studied.** Let  $\ell$  be an odd prime. Fix two absolutely abelian fields  $K_i$  ( $i = 1, 2$ ) which have degree  $\ell$  over  $\mathbb{Q}$  and a prime conductor  $p_i$  ( $p_1 \neq p_2$ ). Note that we necessarily have  $p_i \equiv 1 \pmod{\ell}$ . Suppose also that  $p_1$  splits in  $K_2$  and  $p_2$  splits in  $K_1$ . Let  $d < 0$  be such that

$$\ell \nmid d, \quad \left(\frac{d}{p_i}\right) = 1, \quad \text{and } d \equiv 0, 1 \pmod{4}.$$

Let  $F$  be the imaginary quadratic field  $\mathbb{Q}(\sqrt{d})$  and set  $K = FK_1K_2$ . For future use, we remark that  $(\ell, w_K) = 1$ . The field  $K$  is an abelian extension of  $\mathbb{Q}$  with Galois group  $G = \text{Gal}(K/\mathbb{Q})$  isomorphic to

$$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/\ell\mathbb{Z} \times \mathbb{Z}/\ell\mathbb{Z}.$$

Corollary 3.22 and Theorem 3.23 of [15] imply that there are  $\ell + 1$  minimal cocyclic subgroups which are precisely the subgroups of order  $\ell$  of  $G$  (see Section 3.3 of [15] for the notion of minimal cocyclic subgroup and its connection with 1-covers). Let us denote them by  $H_i$  for  $i = 1, \dots, \ell + 1$  in such a way that  $H_i = G_{p_i}$ , the decomposition group of  $p_i$ , for  $i = 1, 2$ . In order to have a 1-cover, we need to take unramified primes  $p_3, \dots, p_{\ell+1}$  satisfying

$$H_i = \langle \sigma_{p_i} \rangle, \quad \text{for } i = 3, \dots, \ell + 1.$$

By Theorem 3.15 and 3.16 of [15], we see that

$$S = \{\infty, p \mid d, p_1, \dots, p_{\ell+1}\}$$

is a 1-cover for  $K/\mathbb{Q}$  and

$$S_{min} = \{p_1, p_2, \dots, p_{\ell+1}\}.$$

Note that  $|G_{p_i}| = \ell$  for all  $i = 1, 2, \dots, \ell + 1$ . Here is our main theorem for this family of abelian extensions of  $\mathbb{Q}$ .

**Theorem 3.2.** *For  $p \in S_{min}$ , let  $L = K^{G_p}$  and  $\Gamma = \text{Gal}(L/\mathbb{Q})$ . Let also  $R = S \setminus \{p\}$ . Then Question 2.1 has an affirmative answer. More precisely*

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{\ell} \in \mathbb{Z}[\Gamma],$$

and

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{\ell} \in \text{BrSt}_{\mathbb{Z}[\Gamma]}^{(w_L)}(I_L).$$

**Proof:**

Given any prime  $p \in S_{min}$ , letting  $L = K^{G_p}$  as usual, we remark that  $F \subseteq L$ . Set  $H = H_p = \text{Gal}(L/F)$ ; it is a cyclic group of order  $\ell$ . Given  $\alpha \in \mathbb{Z}_\ell[\Gamma]^-$ , there exists a unique  $\tilde{\alpha} \in \mathbb{Z}_\ell[H]$  such that

$$\alpha = e_- \cdot \tilde{\alpha},$$

where  $e_-$  is the usual idempotent

$$e_- = \frac{1-j}{2} \in \mathbb{Z}_\ell[\Gamma].$$

We obtain in this way an isomorphism of rings

$$\psi : \mathbb{Z}_\ell[\Gamma]^- \longrightarrow \mathbb{Z}_\ell[H], \quad \alpha \mapsto \psi(\alpha) = \tilde{\alpha},$$

which allows us to identify  $\mathbb{Z}_\ell[\Gamma]^-$  with  $\mathbb{Z}_\ell[H]$ . At times, we will denote the image of  $\alpha \in \mathbb{Z}_\ell[\Gamma]^-$  simply by  $\tilde{\alpha}$  rather than  $\psi(\alpha)$ .

Since  $(w_L, \ell) = 1$ , one has

$$\theta_{L/\mathbb{Q}}(0) \in \mathbb{Z}_{(\ell)}[\Gamma]^- \subseteq \mathbb{Z}_\ell[\Gamma]^-.$$

The image of  $\theta_{L/\mathbb{Q}}(0)$  via  $\psi$  will be denoted more simply by  $\tilde{\theta}_{L/\mathbb{Q}}$ .

**The ramified primes  $p_1$  and  $p_2$ .** Both ramified primes  $p_1$  and  $p_2$  are treated in the same way. Let us take  $p_1$  and set as usual  $L = K^{G_{p_1}}$ ,  $R = S \setminus \{p_1\}$ ,  $\Gamma = \text{Gal}(L/k)$ , and  $H = \text{Gal}(L/F)$ . Fix a generator  $h$  of  $H$ . Noting that  $p_2$  ramifies in  $L^+$  and splits in  $F$ , Theorem 6.1 of [4] implies that

$$\tilde{\theta}_{L/\mathbb{Q}} = \alpha \cdot (1 - h),$$

for some  $\alpha \in \mathbb{Z}_\ell[H]$  (denoted by  $\vartheta_0$  in loc. cit.). Since  $S(L/\mathbb{Q}) = S(K/\mathbb{Q}) \setminus \{p_1\}$ , we have

$$\theta_{L/\mathbb{Q},R}(0) = \theta_{L/\mathbb{Q}}(0) \cdot \omega_R,$$

where

$$\omega_R \in I_H^{\ell-1}.$$

Therefore,

$$\omega_R = (1 - h)^{\ell-1} \cdot \beta,$$

for some  $\beta \in \mathbb{Z}[H]$ , again since  $H$  is cyclic. Hence, we have

$$\begin{aligned} w_L \cdot \theta_{L/\mathbb{Q},R}(0) &= w_L \cdot e_- \cdot \alpha \cdot \beta \cdot (1 - h)^\ell \\ &= w_L \cdot e_- \cdot \alpha \cdot \beta \cdot \ell \cdot (1 - h) \cdot \gamma, \text{ for some } \gamma \in \mathbb{Z}[H] \\ &= \ell \cdot w_L \cdot \theta_{L/\mathbb{Q}}(0) \cdot \beta \cdot \gamma. \end{aligned}$$

In the previous chain of equalities, we used the fact that  $(1 - h)^\ell \in \ell \cdot (1 - h) \cdot \mathbb{Z}[H]$  as a simple computation, using the binomial theorem, shows. We conclude that

$$\frac{w_L \cdot \theta_{L/\mathbb{Q},R}(0)}{\ell} = w_L \cdot \theta_{L/\mathbb{Q}}(0) \cdot \beta \cdot \gamma \in \mathbb{Z}[\Gamma],$$

and the classical Brumer-Stark conjecture for the special value  $w_L \cdot \theta_{L/\mathbb{Q}}(0)$  implies that Theorem 3.2 is true for the two ramified primes in  $S_{min}$ .

**The unramified primes  $p_3, \dots, p_{\ell+1}$ .** Again, all unramified primes  $p_i$ , for  $i = 3, \dots, \ell+1$ , are treated in the same way. Fix such an  $i$  and let  $L = K^{G_{p_i}}$ . We already know that

$$\frac{w_L \theta_{L/\mathbb{Q},R}(0)}{\ell} \in \mathbb{Z}[\Gamma],$$

because we are in the situation where  $p \in S_{min}$  is unramified in  $K/\mathbb{Q}$  (see the discussion after Hypothesis 2.2). Because of Corollary 2.5, Proposition 2.6, Proposition 2.7 ( $(\ell, w_L) = 1$ ) and Proposition 2.9, we just have to show that

$$\frac{w_L \theta_{L/\mathbb{Q},R}(0)}{\ell} \in \text{Ann}_{\mathbb{Z}[\Gamma]}(Cl_L\{\ell\}^-).$$

Let  $h$  be a generator of  $H$ . Since both  $p_1$  and  $p_2$  are ramified in  $L^+/\mathbb{Q}$ , but split in  $F$ , Theorem 6.1 of [4] shows that there exists  $\alpha \in \mathbb{Z}_\ell[H]$  (in their notation, we have  $h = \gamma$ ,  $\alpha = \vartheta_0$ ,  $s = 2$ , and our  $H$  is their  $\Gamma$ ) satisfying

$$\tilde{\theta}_{L/\mathbb{Q}} = \alpha \cdot (1 - h)^2,$$

and such that

$$(1) \quad \alpha \cdot (1 - h) \in \text{Ann}_{\mathbb{Z}_\ell[H]}(Cl_L\{\ell\}^-).$$

Now, letting  $R = S \setminus \{p_i\}$ , we have

$$\omega_R \in I_H^{\ell-2}.$$

Write

$$\omega_R = \beta \cdot (1 - h)^{\ell-2},$$

for some  $\beta \in \mathbb{Z}[H]$ . We then have

$$\begin{aligned} w_L \cdot \theta_{L/\mathbb{Q},R}(0) &= w_L \cdot e_- \cdot \alpha \cdot (1 - h)^\ell \cdot \beta \\ &= \ell \cdot w_L \cdot e_- \cdot \alpha \cdot (1 - h) \cdot \gamma \cdot \beta, \text{ for some } \gamma \in \mathbb{Z}[H]. \end{aligned}$$

Therefore,

$$\frac{w_L \cdot \theta_{L/\mathbb{Q},R}(0)}{\ell} = w_L \cdot e_- \cdot \alpha \cdot (1 - h) \cdot \gamma \cdot \beta \in \text{Ann}_{\mathbb{Z}[\Gamma]}(Cl_L\{\ell\}^-).$$

Q.E.D.

Here are some remarks. Let us suppose we are in the situation of an unramified prime  $p \in S_{min}$  in the proof of Theorem 3.2. Given a prime  $\mathfrak{p}$  of  $L$ , the classical Brumer-Stark conjecture gives the existence of a  $\varepsilon_R(\mathfrak{p}) \in L^0 \cap \mathcal{A}(L/\mathbb{Q}, w_L)$  such that

$$\mathfrak{p}^{w_L \theta_{L/\mathbb{Q}, R}(0)} = (\varepsilon_R(\mathfrak{p})).$$

In the case where  $\mathfrak{p}$  is relatively prime to the conductor of  $L$ , the algebraic number  $\varepsilon_R(\mathfrak{p})$  is a product of normalized Gauss sums since our base field is  $\mathbb{Q}$ . Instead of simply using results of [4] as we did, it would be possible to show directly that  $\varepsilon_R(\mathfrak{p})$  is a  $\ell$ -th power of an element in  $L^0 \cap \mathcal{A}(L/\mathbb{Q}, w_L)$  using certain complicated identities in group rings and the norm property satisfied by the Euler system of Gauss sums. In fact, this was Greither and Kučera's original approach to related problems (see [5] for instance).

One could ask the following question regarding Question 2.1: If true, do we get new annihilators or not? In the setting of Theorem 3.2, the classical special value

$$w_L \theta_{L/\mathbb{Q}, R}(0)$$

is always in the  $\ell$ -Stickelberger ideal in the sense of Sinnott. The motivating question of [4] was to understand whether or not the Sinnott Stickelberger ideal tells the whole story about the annihilator of the class group. In other words, is it possible to find annihilators outside the Sinnott Stickelberger ideal? They actually found such examples. Using their remarks in that paper, we now explain that in some cases, the special value

$$\frac{w_L \theta_{L/\mathbb{Q}, R}(0)}{\ell}$$

of Theorem 3.2 is *not* in the  $\ell$ -Sinnott Stickelberger ideal as well.

Let us first recall how the Sinnott Stickelberger ideal is defined (for more details see [11]). If both  $E$  and  $F$  are finite abelian extensions of  $\mathbb{Q}$  such that  $E \subseteq F$  and  $R$  is a commutative ring of characteristic 0 (such as  $\mathbb{Z}, \mathbb{Z}_\ell, \mathbb{Q}$ , etc), then the usual restriction homomorphism defined as

$$\sigma \mapsto \sigma|_E,$$

for  $\sigma \in \text{Gal}(F/\mathbb{Q})$  induces a  $R$ -algebra morphism

$$\text{res}_{F/E} : R[\text{Gal}(F/\mathbb{Q})] \longrightarrow R[\text{Gal}(E/\mathbb{Q})].$$

We also need the corestriction map

$$\text{cor}_{F/E} : R[\text{Gal}(E/\mathbb{Q})] \longrightarrow R[\text{Gal}(F/\mathbb{Q})],$$

defined for  $\sigma \in \text{Gal}(E/\mathbb{Q})$  as

$$\text{cor}_{F/E}(\sigma) = \sum_{\substack{\tau \in \text{Gal}(F/\mathbb{Q}) \\ \tau|_E = \sigma}} \tau.$$

Contrary to  $\text{res}_{F/E}$ , the corestriction map is not a morphism of rings, but it is additive. In other words, it is a morphism of  $R$ -modules.

For  $\mathbb{Q}(\zeta_n)$ , where  $n \not\equiv 2 \pmod{4}$ , we have the classical Stickelberger element

$$\theta_n := \sum_{\substack{t=1 \\ (t,n)=1}}^n \frac{t}{n} \sigma_t^{-1} \in \mathbb{Q}[G_n],$$

where  $G_n$  is the Galois group of  $\mathbb{Q}(\zeta_n)/\mathbb{Q}$ . Given  $E/\mathbb{Q}$  we let

$$N_E := \sum_{\sigma \in \text{Gal}(E/\mathbb{Q})} \sigma.$$

One has

$$\theta_{\mathbb{Q}(\zeta_n)/\mathbb{Q}}(0) = \frac{1}{2} N_{\mathbb{Q}(\zeta_n)} - \theta_n,$$

as a consequence of the knowledge of the value at  $s = 0$  of the Hurwitz zeta function. If  $L$  is a finite abelian extension of  $\mathbb{Q}$  with conductor  $m$ , let

$$\theta_L = \text{res}_{\mathbb{Q}(\zeta_m)/L}(\theta_m).$$

Applying the restriction map to the corresponding equality for  $\mathbb{Q}(\zeta_m)$ , we have

$$\theta_{L/\mathbb{Q}}(0) = \frac{1}{2}[\mathbb{Q}(\zeta_m) : L] \cdot N_L - \theta_L.$$

Let us denote the Galois group of  $L/\mathbb{Q}$  by  $\Gamma$ . Let  $S'_L$  be the submodule of  $\mathbb{Q}[\Gamma]$  generated over  $\mathbb{Z}[\Gamma]$  by the elements of the form

$$\text{cor}_{L/L'}(\theta_{L'}),$$

where  $L'$  runs over all subfields of  $L$ . The Sinnott Stickelberger ideal is defined to be

$$S_L = S'_L \cap \mathbb{Z}[\Gamma].$$

If  $p$  is any prime number, the same construction as above replacing  $\mathbb{Z}$  and  $\mathbb{Q}$  by  $\mathbb{Z}_p$  and  $\mathbb{Q}_p$  respectively gives the  $p$ -Sinnott Stickelberger ideal. We shall use the notation  $S'_{L,p}$  and  $S_{L,p}$ . Note that by construction, one has

$$S_L \subseteq S_{L,p},$$

for all prime numbers  $p$ .

Let us come back to the situation of Theorem 3.2, where  $p \in S_{min}$  is an unramified prime,  $|G_p| = \ell$ , and  $L = K^{G_p}$ . Since  $\ell$  is relatively prime to the conductor of  $L$ , we have  $S'_{L,\ell} = S_{L,\ell}$ . Consider the minus  $\ell$ -Sinnott Stickelberger ideal

$$S_{L,\ell}^- = e_- \cdot S_{L,\ell}.$$

If  $L'$  is a subfield of  $L$  which is real with conductor  $m$ , then

$$\theta_{L'} = \frac{1}{2}[\mathbb{Q}(\zeta_m) : L'] \cdot N_{L'}$$

which lives in the plus part. Hence,  $S_{L,\ell}^-$  is generated over  $\mathbb{Z}_\ell[\Gamma]$  by the elements of the form

$$\text{cor}_{L/L'}(\theta_{L'}),$$

where  $L'$  runs over all totally complex subfields of  $L$ . Let  $\tilde{S}_L$  be the image of  $S_{L,\ell}^-$  under the isomorphism  $\psi$  of the beginning of the proof of Theorem 3.2. The ideal  $\tilde{S}_L$  is generated over  $\mathbb{Z}_\ell[H]$  by the elements

$$\text{cor}_{L/L'}(\tilde{\theta}_{L'}),$$

as  $L'$  runs over all totally complex subfields of  $L$ .

As explained on page 1650 of [4] (see comment (2)), the element  $\alpha \cdot (1 - h)$  of equation (1) is outside the Sinnott Stickelberger ideal of  $L$ . Their reasoning is as follows. They work in the quotient ring  $\mathbb{Z}_\ell[H]/N_H$  which is isomorphic to  $\mathbb{Z}_\ell[\zeta_\ell]$  and hence is a discrete valuation ring. The image of  $\tilde{S}_L$  in this quotient ring is generated over  $\mathbb{Z}_\ell[H]$  by a unique element namely

$$\tilde{\theta}_L = -\tilde{\theta}_{L/\mathbb{Q}}.$$

If  $\alpha \cdot (1 - h)$  were in this image, then it would be a multiple of  $\tilde{\theta}_{L/\mathbb{Q}} = \alpha \cdot (1 - h)^2$ . We would then conclude that  $1 - h$  is a unit in this ring which is not the case (the element  $1 - \zeta_\ell$  is not a unit in  $\mathbb{Z}_\ell[\zeta_\ell]$ ).

For simplicity, let us assume further that  $\ell = 3$ . In this case, it is simple to check that both  $\beta$  and  $\gamma$  in the proof of Theorem 3.2 for unramified primes in  $S_{min}$  are also units once projected onto the ring

$$\mathbb{Z}_3[H]/N_H \simeq \mathbb{Z}_3[\zeta_3].$$

Therefore, we conclude as well that

$$\frac{w_L \cdot \theta_{L/\mathbb{Q},R}(0)}{3} \notin S_{L,3}.$$

As a corollary of Theorem 3.2, we get:

**Corollary 3.3.** *In the same setting as in Theorem 3.2, the extended abelian rank one Stark conjecture is true for  $K/\mathbb{Q}$  and the 1-cover  $S$ .*

**Proof:**

If  $p \in S_{min}$  is a ramified prime, Theorem 3.2 implies that the four conditions of Proposition 4.4 of [15] are true; hence,  $St(K/\mathbb{Q}, S, p)$  has an affirmative answer. If  $p \in S_{min}$  is unramified, Theorem 3.2 implies that Question 4.7 of [15] has an affirmative answer. Hence, by Proposition 4.3 of [15], we can conclude that the extended abelian rank one Stark conjecture is true for  $K/\mathbb{Q}$  and the 1-cover  $S$ .

Q.E.D.

#### 4. CONCLUSION

The goal of this paper was to provide further evidence for the extended abelian rank one Stark conjecture by studying two infinite families of finite abelian extensions of  $\mathbb{Q}$ . Rather than studying the extended abelian rank one Stark conjecture itself, we studied a stronger statement, namely an extension of the Brumer-Stark conjecture (Question 2.1). We believe that this extension of the Brumer-Stark conjecture is interesting in itself. There are some new phenomena proper to Question 2.1. Most notably perhaps, in the case where the statement does not reduce to the classical Brumer-Stark conjecture, we are always in a “non semi-simple” situation as explained in §2.3. Moreover, we saw that if true, the extension of the Brumer-Stark conjecture predicts the existence of annihilators which might be outside the Stickelberger ideal in the sense of Sinnott.

If Question 2.1 is true, it would be interesting to find a proof, in the case where the base field is  $\mathbb{Q}$ , along the classical line of Stickelberger’s theorem.

#### REFERENCES

- [1] David S. Dummit and David R. Hayes. Checking the  $\mathfrak{p}$ -adic Stark conjecture when  $\mathfrak{p}$  is Archimedean. In *Algorithmic number theory (Talence, 1996)*, volume 1122 of *Lecture Notes in Comput. Sci.*, pages 91–97. Springer, Berlin, 1996.
- [2] Caleb J. Emmons and Cristian D. Popescu. Special values of abelian  $L$ -functions at  $s = 0$ . *J. Number Theory*, 129(6):1350–1365, 2009.
- [3] Stefan Erickson. An extension of the first-order Stark conjecture. *Rocky Mountain J. Math.*, 39(3):765–787, 2009.
- [4] Cornelius Greither and Radan Kučera. Annihilators of minus class groups of imaginary abelian fields. *Ann. Inst. Fourier*, 57(5):1623–1653, 2007.
- [5] Cornelius Greither and Radan Kučera. Annihilators for the class group of a cyclic field of prime power degree. *Acta Arith.*, 112(2):177–198, 2004.
- [6] Cornelius Greither, Xavier-François Roblot, and Brett A. Tangedal. The Brumer-Stark conjecture in some families of extensions of specified degree. *Math. Comp.*, 73(245):297–315 (electronic), 2004.
- [7] Benedict H. Gross. On the values of abelian  $L$ -functions at  $s = 0$ . *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 35(1):177–197, 1988.
- [8] David R. Hayes. The refined  $\mathfrak{p}$ -adic abelian Stark conjecture in function fields. *Invent. Math.*, 94(3):505–527, 1988.
- [9] David R. Hayes. The conductors of Eisenstein characters in cyclotomic number fields. *Finite Fields Appl.*, 1(3):278–296, 1995.
- [10] Jonathan W. Sands. Galois groups of exponent two and the Brumer-Stark conjecture. *J. Reine Angew. Math.*, 349:129–135, 1984.
- [11] Warren Sinnott. On the Stickelberger ideal and the circular units of an abelian field. *Invent. Math.*, 62(2):181–234, 1980/81.
- [12] Barry Smith. Non-cyclic class groups and the Brumer-Stark conjecture. *J. Number Theory*, 132(2):348–370, 2012.
- [13] Harold M. Stark.  $L$ -functions at  $s = 1$ . IV. First derivatives at  $s = 0$ . *Adv. in Math.*, 35(3):197–235, 1980.
- [14] John Tate. *Les conjectures de Stark sur les fonctions  $L$  d’Artin en  $s = 0$* , volume 47 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1984. Lecture notes edited by Dominique Bernardi and Norbert Schappacher.
- [15] Daniel Vallières. On a generalization of the rank one Rubin-Stark conjecture. *J. Number Theory*, 132(11):2535–2567, 2012.
- [16] Daniel Vallières. *On a generalization of the rank one Rubin-Stark conjecture*. ProQuest LLC, Ann Arbor, MI, 2011. Thesis (Ph.D.)—University of California, San Diego.